

Anticoagulation in Intracerebral Hemorrhage (ICH) Survivors for Stroke Prevention and Recovery

Status: Recruiting

Eligibility Criteria

Sex: Male or Female

Age Group: 18 years and over

This study is NOT accepting healthy volunteers

Inclusion Criteria:

- diagnosis of Intracerebral hemorrhage (ICH) confirmed by brain CT or MRI - documented atrial fibrillation or atrial flutter - can enter study 14 to 180 days after ICH - women willing to use highly effective birth control

Exclusion Criteria:

- prior ICH within last 12 months - women who are pregnant or breast feeding - allergy to aspirin or apixaban - persistent, uncontrolled systolic blood pressure (≥ 180 mm Hg) - contact study staff for additional exclusion criteria

Conditions & Interventions

Interventions:

Drug: Apixaban, Drug: Aspirin

Conditions:

Brain & Nervous System, Heart & Vascular

Keywords:

Anticoagulation, Atrial Fibrillation (AF), CVA, ICH, Intracerebral Hemorrhage, Stroke

More Information

Description: This study will compare the effects of apixaban to aspirin in patients who have atrial fibrillation and a recent brain hemorrhage to see which is better in preventing strokes and death.

Study Contact: Hooman Kamel, MD - hok9010@med.cornell.edu

Principal Investigator: Oladi Benthoo

Phase: PHASE3

IRB Number: SITE00000694

Thank you for choosing StudyFinder. Please visit <http://studyfinder.umn.edu> to find a Study which is right for you and contact sfinder@umn.edu if you have questions or need assistance.