

A Randomized Controlled Trial of Robotic versus Open Radical Hysterectomy for Cervical Cancer (ROCC trial)

Status: Recruiting

Eligibility Criteria

Sex: Female

Age Group: 18 years and over

This study is NOT accepting healthy volunteers

Inclusion Criteria:

- 18 years or older - confirmed cervical cancer with the histological type of adenocarcinoma, squamous cell carcinoma, or adenosquamous carcinoma (Including glassy cell) - Stage IA2, IB1, IB2 cancer - able to care for self and do light work

Exclusion Criteria:

- unable to have a MRI - history of prior pelvic or abdominal radiotherapy - history of cancer less than 5 years ago with the exception of non-melanoma skin cancer.

Conditions & Interventions

Interventions:

Device: da Vinci, Other: open surgery

Conditions:

Cancer, Women's Health

Keywords:

Cervical Cancer

More Information

Description: This study is to investigate if robotic assisted laparoscopy (small incision surgery), is worse than open surgery (otherwise known as a laparotomy) when performing a radical hysterectomy for cervical cancer. Previous research has been done and shown that patients receiving laparoscopy had an increased risk of recurrence of their cervical cancer. But since that time a lot has been learned and improvements have been made, hence why we are relooking at the differences between the two surgical approaches.

Study Contact: Maria Pecoraro - pecor011@umn.edu

Principal Investigator: Colleen Rivard

Phase: N/A

IRB

Number: SITE00001439

Thank you for choosing StudyFinder. Please visit <http://studyfinder.umn.edu> to find a Study which is right for you and contact sfinder@umn.edu if you have questions or need assistance.