

## Addition of a pectoserratus block to interscalene block in patients undergoing total shoulder replacement.

**Status:** Recruiting

### Eligibility Criteria

**Sex:** Male or Female

**Age Group:** 18 years and over

This study is NOT accepting healthy volunteers

**Inclusion Criteria:**

- 18 years to 85 years old - having a total shoulder arthroplasty or reverse total shoulder arthroplasty

---

**Exclusion Criteria:**

- unable to receive regional anesthesia or nerve block - Non-English speakers - women who are pregnant - currently using opioids and people who have chronic pain

### Conditions & Interventions

**Conditions:**

Bone, Joint & Muscle

**Keywords:**

Clinics and Surgery Center (CSC), Shoulder Replacement, Total Shoulder Arthroplasty

### More Information

**Description:** This research is being done to determine if pectoserratus block with bupivacaine will provide improvement in pain control and decrease opioid use, relative to pectoserratus block with a saline solution, among patients undergoing total shoulder arthroplasty.

**Study Contact:** Sherilyn Landree - landr184@umn.edu

**Principal Investigator:** Jacob Hutchins

**IRB**

**Number:** STUDY00019840

---

Thank you for choosing StudyFinder. Please visit <http://studyfinder.umn.edu> to find a Study which is right for you and contact [sfinder@umn.edu](mailto:sfinder@umn.edu) if you have questions or need assistance.