



A Phase 3 Randomized Controlled Trial of Post-Surgical Stereotactic Radiotherapy (SRT) versus Surgically Targeted Radiation Therapy (STaRT) with Gamma Tile for Treatment of Newly Diagnosed Metastatic Brain Tumors.

Status: Recruiting

# Eligibility Criteria

Sex: Male or Female

**Age Group:** 18 years and over This study is NOT accepting healthy

volunteers

### **Inclusion Criteria:**

- one to four newly diagnosed brain metastases, from an extracranial primary tumor (found on MRI) - planned surgery to remove one lesion is between 2.5 cm and 5.0 cm in size, other lesions must be less than 4.0 cm in size - able to complete an MRI of the head with contrast - fluent in English or Spanish language - additional criteria apply, contact study staff

### **Exclusion Criteria:**

- past radiation or surgical therapy newly diagnosed lesion(s) - more than 4 newly diagnosed metastases on MRI - psychiatric, neurologic disease, injury impacting cognition

### Conditions & Interventions

### Interventions:

Device: Gamma Tile-Surgically Targeted Radiation Therapy (STaRT), Radiation: Stereotactic Radiation Therapy

Conditions:

Brain & Nervous System, Cancer

Keywords:

Brain Cancer, Clinics and Surgery Center (CSC)

## More Information

**Description:** The purpose of this research study is to compare surgical tumor removal followed by stereotactic radiotherapy (SRT) to surgical tumor removal followed by radiation therapy delivered by surgically implanted GammaTilesTM (GT). A GammaTile (GT) is an FDA cleared device used to provide radiation therapy following the removal of a brain tumor. GT are small (2cm x 2cm x 0.4cm) collagen squares/tiles that contain sources of radiation that look like grains of rice. If assigned to the GT study group, the doctor will place tiles containing the radiation sources in the cavity left after surgically removing the brain tumor. They do not need to be removed as the collagen tiles will be absorbed and the radiation sources can be left in place. If assigned to the SRT study group, SRT will take place 3-4 weeks after surgery and uses external beams to deliver radiation to the cavity left after surgically removing the brain tumor.

Study Contact: Lydia Golden - golde406@umn.edu

Principal Investigator: Lindsey Sloan

Phase: PHASE3

IRB

Number: SITE00001150

Thank you for choosing StudyFinder. Please visit http://studyfinder.umn.edu to find a Study which is right for you and contact sfinder@umn.edu if you have questions or need assistance.