

Effects of Pain on Laboratory Drinking Topography and Daily Drinking in People with Chronic Temporomandibular Disorder (TMD) Pain

Status: Recruiting

Eligibility Criteria

Sex: Male or Female

Age Group: 18 years and over

This study is NOT accepting healthy volunteers

Inclusion Criteria:

- 21+ years of age, with a valid ID - Fluent in English - Report drinking ≥ 2 days/week - Women must report ≥ 1 drink/day on average over the last 6 months. - Men must report ≥ 2 drink/day on average over the last 6 months - Diagnosis of TMD, myalgia, TMJ, or a combination - Have a smartphone with Internet

Exclusion Criteria:

- History of any psychotic disorder, current major depression, substance use disorder or current alcohol use disorder - Serious medical illness - Under-controlled hypertension or diabetes or neurological disease - Impaired cognitive function - Women who are pregnant breastfeeding, or intending to become pregnant - Loss of sensation in lower legs or physical inability to complete study tasks - Positive drug screening - Medications that contraindicate alcohol use - Comparison group: Chronic pain & TMD

Conditions & Interventions

Interventions:

Drug: alcohol condition, Other: control condition

Conditions:

Bone, Joint & Muscle, Brain & Nervous System, Mental Health & Addiction

Keywords:

Alcohol, Questionnaires, TMD/ TMJ, Virtual Reality

More Information

Description: This study is about how pain affects drinking behaviors in people with chronic temporomandibular disorder (TMD).

Study Contact: ted STUDY - tedstudy@umn.edu

Principal Investigator: Jeff Boissoneault

Phase: EARLY_PHASE1

IRB Number: STUDY00023834

Thank you for choosing StudyFinder. Please visit <http://studyfinder.umn.edu> to find a Study which is right for you and contact sfinder@umn.edu if you have questions or need assistance.